

**Determination of pseudo-Goldstone boson-photon coupling
by the differential time delay of pulsar signals**

Subhendra Mohanty and S. N. Nayak

Institute of Physics, Bhubaneswar-751005, India.

ABSTRACT

Pseudo-Goldstone bosons couple with photons through a P and T violating interaction of the form $\mathcal{L}_{\mathcal{I}} = g_{a\gamma\gamma} a F \tilde{F}$. Strong magnetic fields in rotating compact stars induce a non-zero $\vec{E} \cdot \vec{B}$ outside the stellar surface which acts as a source for the pseudo-scalar field. Pulsar signals propagating through this pseudo-scalar ‘hair’ suffer a differential time lag between the left and the right circularly polarised modes because of the P and T violating pseudoscalar boson-photon interaction. Determination of this time lag upto a microsecond accuracy can lead to a measurement of (or rule out) pseudo-Goldstone boson-photon coupling upto $g_{a\gamma\gamma} \leq 4.6 \times 10^{-9} \text{ GeV}^{-1}$.

e-mail: mohanty@iopb.ernet.in

e-mail: sn@iopb.ernet.in

Goldstone bosons arise in particle physics models when a global symmetry is spontaneously broken. These include axions [1] which arise in many different versions of the Peccei Quinn symmetry breaking and may be the solution of the strong CP problem. Other examples include Majorons [2] which arise when the lepton number is broken spontaneously and are essential in order to reconcile possible heavy neutrinos with astrophysical bounds (a neutrino heavier than 25eV and lighter than a few GeV has to decay into lighter neutrinos and Majorons in order not to overclose the universe). Goldstone bosons also arise in superstring theories [3] in the same multiplet as the graviton and the dilaton and it is possible that some of the above mentioned Goldstone bosons are string axions. The coupling of Goldstone bosons with other fields is determined by the energy scale of the global symmetry breaking. Generically pseudo-scalar boson-photon coupling can be written as $\mathcal{L}_{a\gamma\gamma} = g_{a\gamma\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu}$ with $g_{a\gamma\gamma} \sim C/v$ where C is model dependent constant and v is the global symmetry breaking scale. Strong magnetic fields outside compact rotating stars [4] induce a non-zero $\vec{E} \cdot \vec{B}$ density outside the stellar surface which acts as a source for the pseudo-scalar field. Pulsar signals propagating through this pseudo-scalar ‘hair’ suffer a differential time lag between the left and right circularly polarised modes because of the Parity and Time-reversal violating pseudo-scalar photon interaction. We show that for a typical millisecond pulsar the time lag is $\delta t \simeq 10^{-6}\text{sec}$ for a coupling of the order $g_{a\gamma\gamma} \simeq 10^{-9}\text{GeV}$. Measurement of such a time lag between the two circularly polarised modes of the order of 10^{-6}sec have been reported in literature [5,6,7]. Due to the large systematic error involved these numbers should be regarded as an upper bound on δt which translates to an upper bound on the pseudo-Goldstone boson-photon coupling, $g_{a\gamma\gamma} \leq 4.6 \times 10^{-9}\text{GeV}^{-1}$. Our bound is strictly valid for

massless pseudo-scalar fields. We have estimated that this bound continues to hold for pseudo-scalars with inverse mass greater than the radius of the neutron star i.e. for $m_a < 10^{-10}eV$. Hadronic axions have a non-zero mass (induced by QCD instantons) in the range of $10^{-5} - 10^{-3}eV$, and our bound does not hold for them. Previous astrophysical bounds have considered the coupling of axions to matter in the cooling rates of the Sun, the red giant stars and the Supernova 1987A [8,9,10]. The best astrophysical bound for the symmetry breaking scale as obtained from the cooling rate of red giant stars is $v < 10^{-8}GeV^{-1}$. Terrestrial experiments have attempted to produce axions by laser beams in a cavity with magnetic field. The upper bound on axion-photon coupling from such experiments is $g_{a\gamma\gamma} < 2.5 \times 10^{-6}GeV^{-1}$ for $m_a < 10^{-4}eV$ [11]. Experimental searches for axions in vicinity of the Sun depend on the conversion of axions to X-rays in the stellar magnetic field. Based on a survey of the X-ray flux from the Sun, Lazarus et al [12] give an upper bound of $g_{a\gamma\gamma} < 3.6 \times 10^{-9}GeV^{-1}$ for $m_a < 0.03eV$. Our bound for $g_{a\gamma\gamma}$ is of same order as that of Lazarus et al. In pulsar signals the time lag between the two circularly polarised modes for the current upper bound $g_{a\gamma\gamma} \sim 10^{-9}GeV^{-1}$ is $\delta t \sim 10^{-6}sec$. If it is possible to analyse the pulse profiles at a resolution higher than $10^{-6}sec$ then it may be possible to actually observe a pulse splitting between the left and the right polarised modes and thereby determine the value of $g_{a\gamma\gamma}$. The time lag also depends upon the frequency ($\delta t \propto \omega^{-2}$) and by analyzing pulsar signals at frequencies at lower than $100 MHz$ (for which $\delta t(100 MHz) = 10^{-6}sec$) it may be possible to lower the bounds on $g_{a\gamma\gamma}$ or make a positive measurement of pseudoscalar-photon coupling.

Interaction of pseudo-Goldstone bosons with photons is described by the La-

grangian

$$\mathcal{L} = \frac{1}{2}(\partial_\mu a)(\partial^\mu a) - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}g_{a\gamma\gamma}aF_{\mu\nu}\tilde{F}^{\mu\nu} \quad (1)$$

where $\tilde{F}^{\mu\nu} = (1/\sqrt{-g})\epsilon^{\mu\nu\rho\sigma}F_{\rho\sigma}$ is the dual of $F_{\mu\nu}$. The equation of motion for axion field given by (1) is

$$\left(\frac{1}{\sqrt{-g}}\right)\partial_\mu(\sqrt{-g}\partial^\mu a) = -\frac{g_{a\gamma\gamma}}{4}F_{\mu\nu}\tilde{F}^{\mu\nu}. \quad (2)$$

In terms of the electromagnetic field strengths $E_i \equiv F_{0i}$ and $B^i = \epsilon^{ijk}F_{jk}$, the source term in the right hand side of the equation (2) is $g_{a\gamma\gamma}(1/\sqrt{-g})(\vec{E} \cdot \vec{B})$. In order to have an non-trivial axion field configuration outside the compact stars we must have an external non-zero pseudoscalar density $\vec{E} \cdot \vec{B}$. In dyonic and Kerr-Newmann family of blackholes [13] the external $\vec{E} \cdot \vec{B}$ is generated by the electric/magnetic charges and rotation and acts as a source for the pseudo-scalar hair in black holes [13]. Pulsars have a strong dipolar magnetic field with field strengths $B_0 \simeq 10^{12}$ Gauss which is determined by measuring the frequency of synchrotron X-ray emissions from their neighbourhood. In the oblique rotor model [14] for pulsar emission the dipole moment $\vec{\mu}$ of a pulsar is not aligned along the rotation axis $\vec{\Omega}$ (while in the case of blackhole they are parallel). The dipole axis precesses around the spin axis with an angular frequency Ω and radiates energy at the rate $\dot{E} = (-2/3) |\ddot{\vec{\mu}}|^2 = -(1/6)B_0^2 R^6 \Omega^4 \sin^2\alpha \simeq 10^{38} \text{erg sec}^{-1}$ for a typical pulsar. The angle α can be determined by measuring Ω and $\dot{\Omega}$ for a given pulsar. In the aligned rotor model [14] the angle $\alpha \sim 0$ (a small misalignment is necessary to produce the pulsation) and rate of radiation $\dot{E} \sim -B_0^2 R^6 \Omega^4$. The synchrotron radiation from the plasma accreting along the magnetic poles appears as the pulsed signal to an observer in the cone swept out by the radio beam as it precesses about the spin axis. The instantaneous magnetic field at point $\vec{r}$ is

given by

$$\vec{B} = \frac{B_0 R^3}{r^3} (3\hat{r}(\hat{r} \cdot \hat{\mu}) - \hat{\mu}) \quad (3)$$

where $|\vec{\mu}| = B_0 R^3$ and B_0 is the maximum magnetic field strength at the surface ($r = R$) of the pulsar. Typically $B_0 \simeq 10^8 - 10^{12}$ Gauss and $R \sim 12$ km. Magnetic field in a rotating frame appears as an electric field

$$\vec{E} = -\vec{\Omega} \times \vec{r} \times \vec{B}. \quad (4)$$

The magnetic field (and consequently the electric field) has a time dependent part oscillating with frequency Ω generated by $\vec{\mu}_\perp$, the component of $\vec{\mu}$ perpendicular to $\vec{\Omega}$ which precesses with angular frequency Ω . The time independent part of $\vec{B}$ and $\vec{E}$ are generated by the parallel component $\vec{\mu}_\parallel$ of $\vec{\mu}$. The time average of field strengths at time scale $\geq \Omega^{-1}$ is determined by $\langle \vec{\mu} \rangle = \mu_\parallel = B_0 R^3 \cos \alpha$ (where α is the angle between $\vec{\mu}$ and $\vec{\Omega}$). The average value of electric field outside the stellar surface is then determined by (3) and (4) by matching the boundary conditions at $r = R$. Since $\vec{E} = 0$ in the highly conducting interior region, the component of $\vec{E}$ parallel to the surface is zero at the boundary and the perpendicular component of $\vec{E}$ changes discontinuously at $r = R$. The average electric field outside the stellar surface ($r > R$) is given by [15]

$$\langle \vec{E}^{ext}(r) \rangle = \frac{B_0 R^5 \Omega \cos \alpha}{r^4} \left[3 \left(\sin^2 \theta - \frac{2}{3} \right) \hat{r} - 2 \sin \theta \cos \theta \hat{\theta} \right]. \quad (5)$$

The average value of the pseudoscalar density $\vec{E}^{ext} \cdot \vec{B}$ is

$$\langle \vec{E}^{ext} \cdot \vec{B} \rangle = -B_0^2 \Omega \cos \alpha \frac{R^8}{r^7} \cos^3 \theta \quad (6)$$

which appears as the source term for the pseudo-scalar fields in equation (2). In the absence of pseudo-scalar coupling the strong electric field (5) which is of the

order $E \sim 10^8$ volts/cm (for $\Omega = 1 \text{ sec}^{-1}$ and $B_0 = 10^{12}$ Gauss) will create a space charge of particles with charge density $\rho = (1/4\pi) \vec{\nabla} \cdot \vec{E} = -(1/\pi) \vec{B} \cdot \vec{\Omega} = (B_0 R^3 \Omega / 2\pi r^3)(\sin^2 \theta - 2\cos^2 \theta)$. This space charge is of the order of 10^{11} cm^{-3} and could effectively screen the electric field. Because of the $g_{a\gamma\gamma} a F \tilde{F}$ coupling however the electric field is screened by the pseudo-scalar field and not primarily by the ionised plasma. This follows from the equation of motion of electromagnetic fields modified by the pseudo-scalar coupling term

$$D_\mu F^{\mu\nu} = g_{a\gamma\gamma} (\partial_\mu a) \tilde{F}^{\mu\nu}, \quad (7)$$

which in terms of $\vec{E}$ and $\vec{B}$ gives the modified Maxwells equations

$$\vec{\nabla} \cdot \vec{E} = -g_{a\gamma\gamma} (\vec{\nabla} a) \cdot \vec{B}, \quad (8a)$$

$$-\partial_0 \vec{E} + \vec{\nabla} \times \vec{B} = g_{a\gamma\gamma} [(\vec{\nabla} a) \times \vec{E} + (\partial_0 a) \vec{B}]. \quad (8b)$$

The remaining two Maxwells equations follow from the Bianchi identity $\partial_\mu \tilde{F}^{\mu\nu} = 0$ which gives

$$\vec{\nabla} \cdot \vec{B} = 0, \quad (9a)$$

$$\partial_0 \vec{B} + \vec{\nabla} \times \vec{E} = 0. \quad (9b)$$

From equation (8a) it follows that the effective field experienced by the plasma is

$$\vec{E}_{eff} = \vec{E} + g_{a\gamma\gamma} a \vec{B}. \quad (10)$$

Therefore a nonzero a and $\vec{B}$ will anti-screen the electric field. The space charge induced in the plasma will neutralise E_{eff} in (10) but even with the plasma polarisation $\vec{E}$ can remain non-zero. The standard picture of pulsar magnetosphere [15] is no longer valid in the presence of pseudo-scalar fields. This may have other observable consequences besides the one we study here.

We assume the geometry outside the surface of the pulsar is described by the Kerr metric

$$ds^2 = \left(1 - \frac{2GM}{r}\right) dt^2 - \left(1 - \frac{2GM}{r}\right)^{-1} dr^2 - r^2 (d\theta^2 + \sin^2\theta d\phi^2) - \frac{4GMA}{r} \sin^2\theta d\phi dt. \quad (11)$$

The form (11) is an approximation to the Kerr metric, valid as long as the angular momentum per unit mass $A \ll GM$. For a typical millisecond pulsar ($M = M_\odot, R = 12 \text{ km}$), we have $A = 10^{-6} \text{ km}$ and $GM = 1.5 \text{ km}$. Therefore (11) is an adequate approximation to the Kerr metric. For the metric given by (11) the equations of motion (2) for the axisymmetric, time independent solution $a(r, \theta)$ reduces to

$$\begin{aligned} \frac{1}{r^2} \frac{\partial}{\partial r} \left[r^2 \left(1 - \frac{2GM}{r}\right) \frac{\partial}{\partial r} a(r, \theta) \right] + \frac{1}{r^2 \sin^2\theta} \frac{\partial}{\partial \theta} \left[\sin\theta \frac{\partial}{\partial \theta} a(r, \theta) \right] \\ = g_{a\gamma\gamma} (B_0^2 R^8 \Omega \cos\alpha) \frac{1}{r^7} \cos^3\theta. \end{aligned} \quad (12)$$

The Greens function which is solution of $\nabla^2 G(x, y) = -\delta^3(x - y)/\sqrt{-g}$ for the Laplacian operator in (12) is given by [13]

$$\begin{aligned} G(r, \theta, r_0, \theta_0) &= \sum_{l=0}^{\infty} C_l P_l \left(\frac{r_0}{GM} - 1 \right) Q_l \left(\frac{r}{GM} - 1 \right) P_l(\lambda), \quad r > r_0 \\ &= \sum_{l=0}^{\infty} C_l Q_l \left(\frac{r_0}{GM} - 1 \right) P_l \left(\frac{r}{GM} - 1 \right) P_l(\lambda), \quad r < r_0, \end{aligned} \quad (13)$$

where $\lambda = \cos\theta \cos\theta_0 + \sin\theta \sin\theta_0$, $C_l = (2l - 1)/4\pi M$ and P_l and Q_l are the two associated Legendre functions. The solution of the inhomogenous equation (12) is

$$\begin{aligned} a(r, \theta) &= - \int_R^r dr_0 \int_0^\pi d\theta_0 \int_0^{2\pi} d\phi_0 r_0^2 \sin^2\theta_0 G(r, \theta, r_0, \theta_0) \\ &\quad \times \left(g_{a\gamma\gamma} B_0^2 R^8 \cos\alpha \Omega \right) \frac{\cos^3\theta_0}{r_0^7}. \end{aligned} \quad (14)$$

Substituting (13) in (14) and evaluating the integral we obtain the Goldstone boson configuration outside pulsar ($r > R$)

$$a(r, \theta) = -\left(\frac{2}{575}\right) \frac{g_{a\gamma\gamma} B_0^2 R^8 \Omega \cos\alpha}{(GM)^3} \frac{\cos\theta}{r^2} + O\left(\frac{1}{r^3}\right). \quad (15)$$

Propagation of electromagnetic radiation through this coherent pseudo-Goldstone boson field is governed by the wave equation obtained by combining the Maxwells equations (8) and (9)

$$\nabla_\mu \nabla^\mu \vec{B} = -g_{a\gamma\gamma} (\vec{\nabla} a) \times \partial_0 \vec{B}. \quad (16)$$

With the eikonal ansatz

$$\vec{B}(x, t) = \vec{\mathcal{B}} e^{iS(x, t)} \quad (17a)$$

and

$$-i\partial_\mu \vec{B} = (\partial_\mu S) \vec{B} = k_\mu \vec{B} \quad (17b)$$

the wave equation reduces to

$$k_\mu k^\mu \vec{\mathcal{B}} = i g_{a\gamma\gamma} (\vec{\nabla} a \times \omega \vec{\mathcal{B}}) \quad (18)$$

where $k^\mu = (\omega, \vec{k})$ is the photon four momentum and $\vec{\mathcal{B}}$ is the magnetic field of the radiation. The pulsed radiation from pulsars originates from the polar regions of the pulsar [16]. In this region $a \simeq a_0(1/r^2)$ and $\vec{\nabla} a = -(2a_0/r^3) \hat{r}$ is radial. Consider right and left circularly polarised combinations for the transverse magnetic field $\vec{\mathcal{B}}$

$$\vec{\mathcal{B}}_\pm = (\vec{\mathcal{B}}_i \pm i\vec{\mathcal{B}}_j) \quad (19)$$

where $\vec{\mathcal{B}}_i$ and $\vec{\mathcal{B}}_j$ are the components of $\vec{\mathcal{B}}$ along any two fixed directions $\hat{e}_i$ and $\hat{e}_j$ orthogonal to $\hat{r}$. Substituting (19) in (18) we find that the equations for the two circularly polarised modes $\vec{\mathcal{B}}_\pm$ decouple as

$$k_\mu k^\mu \vec{\mathcal{B}}_\pm \mp g_{a\gamma\gamma} (\partial_r a) \omega \vec{\mathcal{B}}_\pm = 0. \quad (20)$$

This shows that the propagating modes in a background with non-zero pseudoscalar gradients are circularly polarised modes [17] with different phase velocities. The dispersion relation for the $\vec{B}_\pm$ modes propagating radially from the poles is

$$\omega^2 \left(1 - \frac{2GM}{r}\right)^{-1} - k_r^2 \left(1 - \frac{2GM}{r}\right) = \pm g_{a\gamma\gamma} (\partial_r a) k_0. \quad (21)$$

The Eikonal phase can be written as

$$S(r, t) = \omega t - \int_R^r k_r dr. \quad (22)$$

The propagation time t to reach a point r is given by the orbit equation $\partial S / \partial \omega = 0$ which gives

$$t = \frac{\partial}{\partial \omega} \int_R^r k_r dr. \quad (23)$$

Substituting for k_r from (20) we have

$$t_\pm = \int_R^r dr \left(1 + \frac{2GM}{r}\right) \mp g_{a\gamma\gamma}^2 \int_R^r dr \frac{(\partial_r a)^2}{4\omega^2} \left(1 - \frac{2GM}{r}\right), \quad (24)$$

where $t_+(t_-)$ are the arrival times of the $+$ ($-$) circularly polarised modes. Substituting for $\partial_r a(r)$ from (15) (with $\cos\theta = 1$) and evaluating the integrals in (24) we obtain the arrival times of the pulses from a pulsar at distance r from the earth

$$t_\pm = r + 2GM \ln\left(\frac{r}{R}\right) \pm \frac{1}{5} \left(\frac{2}{575}\right)^2 \frac{g_{a\gamma\gamma}^4 B_0^4 R^{11} \Omega^2 \cos^2 \alpha}{\omega^2 (GM)^6}. \quad (25)$$

The second term is the familiar time delay due to the gravitational field of the pulsar and the last term shows the birefringence properties of the pseudo-scalar background. The differential time delay between the positive and the negative polarised modes is

$$\delta t = t_+ - t_- = \frac{2}{5} \left(\frac{2}{575}\right)^2 \frac{g_{a\gamma\gamma}^4 B_0^4 R^{11} \Omega^2 \cos^2 \alpha}{\omega^2 (GM)^6}. \quad (26)$$

Pulsar signals are recorded in the radio frequency range as low as $\omega \simeq 100MHz$. For a typical millisecond pulsar $M = M_\odot = 10^{57}GeV$, $B_0 = 10^{12}Gauss \simeq 10^{-9}GeV^2$, $R = 12 km = 6 \times 10^{19}GeV^{-1}$, $\cos\alpha \simeq 1$ the differential time delay turns out to be 10^{-6} sec for $g_{a\gamma\gamma} \simeq 10^{-9}GeV^{-1}$. By putting an experimental bound on δt one can therefore obtain the following upper bound on the pseudo-Goldstone boson-photon coupling.

$$g_{a\gamma\gamma} = 4.6 \times 10^{-9}GeV^{-1} \left(\frac{\delta t}{10^{-6}sec} \right)^{1/4} \left(\frac{10^{12}G}{B_0} \right) \left(\frac{12km}{R} \right)^{11/4} \left(\frac{10^{-3}sec}{\Omega} \right)^{1/2} \times \left(\frac{\omega}{100MHz} \right)^{1/2} \left(\frac{M}{M_\odot} \right)^{3/2}. \quad (27)$$

Loseco et al [5] have analysed the signals from the millisecond pulsar PSR 1937+214 using the data of Cordes and Stinebring [6]. They report that a time lag 5×10^{-6} sec is observed between the positive and the negative polarised modes. Klien and Thorsett [7] report a differential time lag of $0.37 \pm 0.67\mu sec$. At the one sigma limit one can put an upper bound on the time lag to be $\delta t \leq 10^{-6}$ sec. This leads to the upper bound on the goldstone boson photon coupling

$$g_{a\gamma\gamma} \leq 4.6 \times 10^{-9}GeV^{-1}. \quad (28)$$

From (26) it is clear that by going for lower frequency pulses one should observe a larger time lag since $\delta t \propto \omega^{-2}$. Therefore by analysing lower frequency signals it should be possible to observe a non-zero time lag which would be a positive signature of pseudo-scalar photon interaction.

References

- 1 R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38** (1977) 1440; Phys. Rev. **D16** (1977) 1791.
- 2 Y. Chikashige, R. N. Mohapatra and R. D. Peccei, Phys. Rev. Lett. **45** (1980) 1926; Phys. Lett **B 98** (1981) 265;
G. B. Gelmini and M. Roncadelli, Phys. Lett **B 99** (1981) 411;
H. Georgi, S. L. Glashow and S. Nussikov, Nucl. Phys. **B 193** (1981) 297.
- 3 M. Green, J. Schwarz and E. Witten, ‘Superstring theory’ Cambridge U.P; Cambridge, 1987.
- 4 S. L. Shapiro and S. A. Teukolsky, ‘Black holes, White dwarfs, and Neutron stars’, John Wiley and Sons, (1983).
- 5 J. M. Loseco, G. E. A. Matsas, A. A. Natale and J. A. F. Pacheco, Phys. Lett **A 138** (1989) 5.
- 6 J. M. Cordes and D. R. Stinebring, Ap J. Lett. **277** (1984) 53.
- 7 J. R. Klien and S. E. Thorsett, Phys. Lett **A 145** (1990) 79.
- 8 G. G. Raffelt, Phys Rep. **198** (1990) 1.
- 9 M. S. Turner, Phys Rep. **197** (1990) 67.
- 10 E. W. Kolb and M. S. Turner, ‘The Early Universe’, Addison-Wesley, (1990).
- 11 Y. Semertzidis et al, Phys. Rev. Lett. **25** (1990) 2988.
- 12 D. M. Lazarus et al, Phys. Rev. Lett. **69** (1992) 2333.
- 13 B. Campbell, N. Kaloper and K. A. Olive, Phys. Lett **B 263** (1991) 364;
Phys. Lett **B 285** (1992) 199.
- 14 F. Pacini, Nature **216** (1967) 567; Nature **219** (1968) 145;
J. E. Gunn and J. P. Ostriker, Nature **221** (1968) 254.
- 15 P. Goldreich and W. H. Julian, Ap. J. **157** (1969) 869.

- 16 V. Radhakrishnan and D. J. Cooke, *Astrophys. Lett.* **3** (1969) 225.
- 17 Sean M. Carroll, George B. Field and Roman Jackiw, *Phys. Rev.* **D41** (1990) 1231;
Sean M. Carroll and George B. Field, *Phys. Rev.* **D43** (1991) 3789.